

## Prestop® For Effective Biological Control Of Botrytis On Ornamentals, Herbs & Vegetables

### PRESTOP®

- EPA Registered (EPA 64137-11) and OMRI listed biofungicide
- Based on a naturally occurring soil fungus *Gliocladium catenulatum* strain J1446
- Easy and safe to use
- Compatible with IPM programs
- Safe to beneficials
- Good resistance management tool for rotation of fungicides
- No PHI is needed
- Package size 3.5 oz. (100 gr.) and 2.2 lb (1 kg.)

# PRESTOP®

# Verdera



### Prestop® defence against *Botrytis*

Prestop® offers an effective tool to protect your crop from this devastating pathogen. Prestop® is an environmentally friendly biofungicide that is exceptionally safe for both users and plants. Prestop® works best as a preventative fungicide. First treatments applied at the seedling stage or at potting/planting are ideal. The active substance is a living microbe, which survives on foliage for several weeks protecting the crop against the invading pathogen.

### How to use PRESTOP®


**Rate** Use 0.5 % aqueous suspension for spraying (example - 100 gr. (3.5 oz.) of PRESTOP®/20 L (5.2 gallons) of water). Spray to wet with sufficient volume of solution ensuring good coverage of plants.

**Timing** Start applications at seedling stage or at potting/planting. Repeat the treatment at the intervals of 3 to 4 weeks.

### Preparing aqueous PRESTOP® suspension

To prepare 0.5% *suspension*, mix 100 gr. (3.5 oz.) of PRESTOP® carefully in a small volume of water (e.g. 2 L (2 quarts)). Agitate as needed to get the product evenly dispersed before diluting to final volume of 20 L (5.2 gallons). The water *suspension* should be used within 24 hours of being prepared if stored at room temperature. The *suspension* keeps for 7 days at temperatures below +8 °C (46 °F).

Effect of Prestop® on disease severity of *Botrytis* on pelargonium (ELMHIRST 2006, Canada)



Another Way  
We Are  
Helping You Grow

# PRESTOP® on Ornamental Plants


## PRESTOP®

- Effectively prevent *Pythium*, *Phytophthora*, *Rhizoctonia* on ornamental plants.
- Incorporate PRESTOP® on your mixing line or as a drench via irrigation system.
- Added insurance--dip cuttings into a 0.5% aqueous PRESTOP® suspension before planting.
- Resistance of pathogens to Prestop® are unlikely to occur.

## How to use PRESTOP®

**Rate** Use 0.5 % aqueous suspension for spraying (example - 100 gr. (3.5 oz.) of PRESTOP®/20 L (5.2 gallons) of water). Spray to wet with sufficient volume of solution ensuring good coverage of plants.

**Timing** Start applications at seedling stage or at potting/planting. Repeat the treatment at the intervals of 3 to 4 weeks.



Do not treat woody, unrooted cuttings.